

LA-UR-80-1611

4

TITLE: Metal Sulfite/Sulfate Reactions in Thermochemical Hydrogen Cycles\*

**MASTER**AUTHOR(S): C.F.V. Mason, CMB-3  
M. G. Bowman, CMB-D0**DISCLAIMER**

This document contains information developed by an agency of the United States Government. The United States Government and its agency, the Los Alamos Scientific Laboratory, make no warranty, expressed or implied, for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily imply its endorsement, recommendation, or favor by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

SUBMITTED TO: To be presented and included in the Proceedings of the 3rd I.E.A. Thermochemical Hydrogen Workshop held at the Tsukuba Center for Science and Technology, Japan, June 18-21, 1980.

University of California

\*Work completed under the auspices of the U.S. Department of Energy.  
Office of Basic Energy Sciences, Division of Chemical Sciences.

By acceptance of this article, the publisher recognizes that the U.S. Government retains a nonexclusive, royalty free license to publish or reproduce the published form of this contribution, or to allow others to do so, for U.S. Government purposes.

The Los Alamos Scientific Laboratory requests that the publisher identify this article as work performed under the auspices of the U.S. Department of Energy

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

**LOS ALAMOS SCIENTIFIC LABORATORY**

Post Office Box 1663 Los Alamos, New Mexico 87545  
An Affirmative Action/Equal Opportunity Employer



## Metal Sulfite/Sulfate Reactions in Thermochemical Hydrogen Cycles\*

C.F.V. Mason and M. G. Bowman


Los Alamos Scientific Laboratory,  
Los Alamos, New Mexico 87545

The thermochemical cycles which have been most extensively developed at this time<sup>1,2,3</sup> all involve the thermal decomposition of sulfuric acid which is corrosive. Metal sulfate cycles have been studied<sup>4,5</sup> as a means of circumventing handling corrosive mixtures at high temperatures. However, these metal sulfate cycles still use an electrochemical step to produce H<sub>2</sub>. This abstract examines alternate H<sub>2</sub> producing steps to be used in conjunction with metal sulfates.


The simplest example of such a reaction involves direct oxidation of a metal sulfite by water:



However, although conceptually possible, when experimentally investigated<sup>6</sup> no reaction was obtained. A modification was tried using magnetite:



$$\Delta G_0^\circ = 540 \text{ K}$$




Again, no reaction was observed.

Another type of sulfite oxidation, for which experimental proof has been obtained, utilizes I<sub>2</sub>:

---

\*Work completed under the auspices of the U.S. Department of Energy, Office of Basic Energy Sciences, Division of Chemical Sciences.




$$\text{where } \text{M} = \text{Mg}, \Delta H_{298}^\circ = -333.1 \text{ kJ} \quad (5\text{a})$$

$$\text{M} = \text{Ca}, \Delta H_{298}^\circ = -255.7 \text{ kJ} \quad (5\text{b})$$

$$\text{M} = \text{Ba}, \Delta H_{298}^\circ = -265.7 \text{ kJ} \quad (5\text{c})$$

when  $\text{M} = \text{Ca}$  or  $\text{Ba}$ , product separation is trivial since their sulfates are insoluble. However,  $\text{MgSO}_4$  is soluble so for reaction (5a) the problem of separating sulfate and iodide remains. Drying, hydrolysing  $\text{MgI}_2$  and thermally decomposing the resultant HI replaces the electrochemical and drying steps in the Westinghouse cycle. Drying and hydrolysing  $\text{MgI}_2$  replaces drying and separating HI in the GA cycle. Experimentally, when using dilute solution ( $0.2 \text{ mol l}^{-1}$ ) at room temperature all these reactions proceed rapidly and are probably complete within the first minute of mixing (as evidenced from the color change from a grey suspension to a white suspension and brown solution). They have all been shown to be complete within 1 h ( $> 95\%$ ). In case (5a) a brown solution is formed since  $\text{MgSO}_4$  is soluble. In the Ca case, the reaction was repeated using the half hydrate at higher concentration. At  $2.0 \text{ mol l}^{-1}$  ( $\text{H}_2\text{O}/\text{CaSO}_3 = 28$ ), the reaction was complete, whereas at  $4.0 \text{ mol l}^{-1}$  ( $\text{H}_2\text{O}/\text{CaSO}_3 = 14$ ) 73% of the reaction took place. This relatively low yield was due both to the exothermicity which caused some of the  $\text{I}_2$  to vaporize and also to the mud-like nature of the reactants which made stirring difficult.

The cycle is completed by thermally decomposing the sulfate (as well as hydrolysing  $\text{MgI}_2$  and thermally decomposing the resultant HI). Both Ba and Ca sulfates decompose above the temperature range often considered available for such cycles:







$$\text{where } \text{M} = \text{Mg}, \Delta G_{1170}^\circ = 0 \quad (6\text{a})$$

$$\text{M} = \text{Ca}, \Delta G_{1810}^\circ = 0 \quad (6\text{b})$$


$$\text{M} = \text{Ba}, \Delta G_{2030}^\circ = 0 \quad (6\text{c})$$

Conceptually, however, an insoluble sulfate with a lower decomposition temperature can be used.

An extension to this class of reactions was also examined:







The advantages of these reactions is that the oxide lowers the decomposition temperature of the sulfate:<sup>7</sup>



The observed course of reaction (7a) in dilute solution ( $0.2 \text{ mol l}^{-1}$ ) leaves  $\text{CaWO}_4$  unchanged but converts all the  $\text{I}_2$  to iodide. Some  $\text{SO}_2$  is evolved. Reactions (7b,c,d) occur as written,  $\text{Ca}(\text{or Ba})\text{SO}_4$  is precipitated and all the  $\text{I}_2$  is converted to iodide. No oxide has been seen in the x-ray powder diffraction pattern of the solid product, but the weights obtained suggest it is present. Reactions (7b,c,d) were also carried out in acetone in order to cut down the heat of vaporization of the solvent ( $+30.5 \text{ kJ mol}^{-1}$  vs.  $+43.9 \text{ kJ mol}^{-1}$  for water). However, in all cases,  $\text{MXO}_4$  was recovered unchanged at the end of the reaction time (0.5 h). In an allied concentrated aqueous reaction using  $\text{BaWO}_4$  and  $\text{MgSO}_3$  ( $2.9 \text{ mol l}^{-1}$ ),  $\text{BaWO}_4$  was recovered unchanged although  $\text{MgSO}_3$  was completely converted to sulfate. Note that in dilute aqueous solution ( $0.2 \text{ mol l}^{-1}$ ),  $\text{BaWO}_4$  was completely converted to  $\text{BaSO}_4$  and  $\text{WO}_3$ .

This work is being extended to include a similar cycle based on lanthanum:



Unlike reaction (5), reaction (9) is thermodynamically possible in the absence of water. In order to effect a separation of the products, however, a solvent for one of the products must be used.  $\text{La}_2(\text{SO}_4)_3$  is insoluble in water (especially at elevated temperatures) while  $\text{LaI}_3$  is soluble.  $\text{La}_2(\text{SO}_4)_3$  is also insoluble in acetone. Acetone, not only requires less energy for vaporization than water, it also, by allowing the reaction to be carried out in the absence of water, prevents the formation of hydrates.

Reaction (11) has been studied<sup>7</sup> and reasonable rates are obtained at 1075K.  $\text{La}_2(\text{SO}_4)_3$ , itself, decomposes at 1270 K.<sup>8</sup>

#### References

1. G. H. Parker and P.W.T. Lu, Proc. 14th IECEC Meeting, Boston (1979).
2. G. Besenbruch, *ibid.*
3. D. Van Velzen and H. Langenkamp, *ibid.*
4. M. G. Bowman, Proc. 1st National Topical Meeting on Nuclear Process Heat Applications, American Nuclear Soc., Los Alamos (1974).
5. K. E. Cox, W. M. Jones, and C. L. Peterson, Proc. 3rd World  $\text{H}_2$  Energy Conference, Tokyo (1980).
6. C. F. V. Mason, Unpublished Results (1979).
7. C. M. Hollabaugh, T. C. Wallace, M. G. Bowman, and W. M. Jones, Proc. 3rd World  $\text{H}_2$  Energy Conference, Tokyo (1980).
8. M. W. Nathans and W. W. Wendlandt, J. Inorg. Nucl. Chem., 24, 869 (1962).